Fire Points — C3

Joseph Audette, MD


Important Signal of Pathology

- Check Fire points as a routine after Hara Diagnosis
- Majority only positive it tender
- **Exceptions:**
 - Kidney Fire point usually very sensitive if positive, but occasionally in someone with Adrenal Exhaustion and Kidney Deficiency, it will only be tight
 - Spleen Fire point often sensitive given location on first toe near Bunion
 - May or may not be clinically relevant

Yin Meridians Hand: Fire Point Locations

Meridian	FIRE	METAL	WATER
Lung	LU 10 – Japanese location, at thenar eminence in muscle belly of Abductor Pollicis Brevis	LU 8 – Located 1 cun proximal to LU 9 (wrist crease) at the level of the radial styloid process	LU 5 – Japanese location, halfway between the lateral edge of cubital crease and the radial side of the biceps tendon in brachioradialis muscle
Pericardium	PC 8 – Between 2 nd and 3 rd metacarpal bones, closer to radial side of 3 rd , where tip of middle finger falls with loose fist	PC 5 – Japanese location in region of TCM PC 5, which is 3 cun proximal from PC 7 (wrist crease), but can vary proximal and distal to this location based on which point releases PC 8	PC 3 – On Transverse cubital crease on either radial side of biceps tendon (Japanese point) or ulnar side (TCM point)
Heart	HT 8 – Located between the 4 th and 5 th metacarpal bones, where the small finger lands when making a loose fist	HT 4 – Located 1.5 cun proximal to HT 7 (wrist crease), on radial side of the flexor carpi ulnaris tendon	HT 3 – Located at the end of the ulnar side of the cubital crease when the elbow is fully flexed, near medial epicondyle of elbow


Yin Meridians Hand: Fire Point



Yang Meridians Hand: Fire Point Locations

Meridian	FIRE	METAL	WATER
Large Intestine	LI 5 – Located on radial aspect of wrist in anatomic snuffbox, between tendons of extensor pollicis longus and brevis	LI 1 - Located at corner of 2 nd finger nail bed on radial side	LI 2 – Located distal to metacarpal-phalangeal joint of 2 nd finger (TCM) Alternate Japanese location is distal to proximal interphalangeal joint of 2 nd finger
Triple Warmer	TW 6 – Located 3 cun proximal from TW 4 (wrist crease) between radial and ulna bones	TW 1 – Located at corner of 4 th finger nail bed on ulnar side	TW 2 – Located distal to metacarpal-phalangeal joint of 4 th finger, ulnar side
Small Intestine	SI 5 – Located at ulnar border of wrist in the depression between the head of the ulna and triquetral bone	SI 1 – Located at corner of 5 th finger nail bed on ulnar side	SI 2 – Located distal to metacarpal-phalangeal joint of 5 th finger, ulnar side

Yang Meridians Hand: Fire Point Locations





Yin Meridians Foot: Fire Point Locations

Meridian FIRE **METAL WATER** SP 5 - Located in the SP 9 - On the medial Spleen SP 2 - Located on the depression distal, aspect of the lower leg, in medial side of the great the depression of the anterior and inferior toe, anterior and inferior lower border of the to the medial to the 1st metatarsomalleolus medial condyle of the phalangeal joint at the tibia. junction of the red and white skin KD 10 - When the knee is Kidney KD 2 - On the medial KD 7 - On the medial flexed, medial to aspect of the lower aspect of the foot, below leg, 2 cun directly popliteal fossa, between the tuberosity of the above KID 3, anterior the tendons of m. navicular bone - a to Achilles tendon semitendinosus and region that extends to semimembranosus. plantar surface Liver LV 2 - On the dorsum of LV 4 - On the dorsum LV 8 – In the depression on the medial end of the of the foot, between SP foot, proximal to web 5 and ST 41, in the transverse popliteal margin, between the 1st depression on the crease, in depression and 2nd toes distal to medial side of the anterior to the tendons of lateral 1st metatarsal tendon m. tibialis the m. semitendinosus phalangeal joint anterior. and semimembranosus.


Yin Meridians Foot: Fire Point Locations



Yang Meridians Foot: Fire Point Locations

Meridian	FIRE	METAL	WATER		
Stomach	ST 41 – At ankle. in the depression between the tendons m. extensor hallucis longus and digitorum longus.	ST 45 – On the foot, on the lateral side of the end of the second toe, . 0.1 cun from the corner of the nail.	ST 44 – On the dorsum of the foot, proximal to the web margin between the second and third toes, distal to the 2 nd metatarso-phalangeal joint		
Bladder	BL 60 – On the foot, behind the lateral malleolus, in the depression between the tip of the lateral malleolus and Achilles tendon.	BL 67 – On the lateral side of the small toe, 0.1 cun from the corner of the nail.	BL 66 – On the lateral aspect of the foot, distal to the 5th metatarsophalangeal joint, at the junction of the red and white skin.		
Gallbladder	GB 38 – Located 4 cun above the tip of the lateral malleolus, slightly anterior to the anterior border of the fibula.	GB 44 – On the foot, on the lateral side of the 4th toe, 0.1 cun from the corner of the nail.	GB 43 – On the lateral side of the foot, proximal to the margin of the web between the 4th and 5th toes, distal to the 4th metatarso-phalangeal joint		


Yang Meridians Foot: Fire Point Locations



10 Stems: Creation Cycle

03


- Creation Cycle leads natural combinations of Fire Point treatments
 - Metal is Mother of Water
 - Often find Kidney and Lung Fire points active
 - Triple warmer (Fire) and Gallbladder M/W treatments often combined



10 Stems: Control Cycle

CB

- Control cycle also involved with combinations of Fire Points
 - Liver and Large Intestine Fire points often positive
 - Gallbladder and Spleen Fire points can be active together
- Also see Yin/Yang interactions
 - Liver and Gallbladder fire points both active
 - Stomach and Spleen



Summary

OB

- Extremely useful part of our diagnostic evaluation
- Get into habit of checking on every patient
- When there are pairs of fire points positive on different meridians, it may provide clues to issues that are occurring on control or creation cycle
 - For example when Lung and Kidney fire point tender, indicates Lung disturbance affecting ability to nourish Kidney
 - Recommend breathing exercises to patient